

# Notes on the Li & Paczyński “kilonova paper”

Edoardo Milotti

December 19, 2023

In this short handout I provide a quick derivation of some equations in the Li & Paczyński paper [1] (LP).

- LP denote the speed with  $V$ , but we also need a symbol for volume, so here I use  $v$  for speed and  $V$  for volume.
- Here I use the standard convention that  $U$  is the energy and  $u$  is the energy density.
- LP denote the entropy per unit mass with the symbol  $S$ ; following the standard conventions I use  $S$  for entropy and  $S/M$  for the entropy per unit mass.
- From the equations for the energy density of blackbody radiation it can be shown that the energy density is

$$u = 3P = aT^4 \quad (1)$$

where  $P$  is the pressure,  $T$  is the temperature of the photon gas, and

$$a = \frac{8\pi^5 k_B^4}{15h^3 c^3} \approx 7.56 \times 10^{-16} \text{ J K}^{-4} \text{ m}^{-3}. \quad (2)$$

The parameter  $a$  is related to the Stefan-Boltzmann constant:  $\sigma = ac/4$ .

It can also be shown that the entropy is

$$S = \frac{4}{3}aVT^3 \quad (3)$$

Then, the internal energy change per unit mass in the LP paper (equation 3 in the paper) is

$$\frac{TdS}{M} = \frac{dU}{M} = d\left(\frac{U}{M/V}\right) = d\left(\frac{u}{\rho}\right) = \frac{du}{\rho} + u d\left(\frac{1}{\rho}\right). \quad (4)$$

Using equation 1 in the paper

$$\rho = \frac{3M}{4\pi v^3} t^{-3}, \quad (5)$$

we find

$$\frac{dU}{M} = \frac{4\pi v^3}{3M} (t^3 du + 3ut^2 dt). \quad (6)$$

- Equation 5 for radioactive heating in LP, is obtained as follows, using the rate  $\lambda_{\text{rad}} = t_{\text{rad}}^{-1}$  instead of  $t_{\text{rad}}$ . A uniform distribution of  $\ln t_{\text{rad}}$  corresponds to a power law distribution of  $t$  which which corresponds to the probability

$$\sim t^{-1} dt = \lambda \left| d \left( \frac{1}{\lambda} \right) \right| = \frac{d\lambda}{\lambda} = d \ln \lambda, \quad (7)$$

i.e., we obtain a uniform distribution in  $\ln \lambda$ . Then we have to compute the integral

$$f c^2 \int_{\lambda_{\min}}^{\lambda_{\max}} \lambda e^{-\lambda t} \frac{d\lambda}{\lambda} = - \frac{f c^2}{t} e^{-\lambda t} \Big|_{\lambda_{\min}}^{\lambda_{\max}} \approx \frac{f c^2}{t}. \quad (8)$$

- Radiative losses. Substituting the expressions for  $u$ ,  $R = vt$  and  $\sigma$  in the expression  $L = 4\pi R^2 F$  we find the heat loss

$$L = 4\pi R^2 F = 4\pi R^2 \frac{\sigma T^4}{\kappa \rho R} = 4\pi R^2 \frac{acT^4}{4\kappa R} \frac{4\pi v^3 t^3}{3M} = \frac{4\pi^2 c}{3\kappa M} (aT^4)(v^4 t^4) = \frac{4\pi^2 c v^4}{3\kappa M} ut^4, \quad (9)$$

which is equation 6 in LP.

- If  $\kappa \rho R = 1$ , then we obtain equation 7 in LP as follows:

$$R = vt_c = \frac{1}{\kappa \rho} = \frac{4\pi v^3 t_c^3}{3\kappa M}, \quad (10)$$

i.e.,

$$t_c = \left( \frac{3\kappa M}{4\pi v^3} \right)^{1/2}. \quad (11)$$

- Starting from the heat balance

$$\frac{L}{M} = \epsilon - T \frac{d(S/M)}{dt}, \quad (12)$$

we find

$$\frac{1}{M} \left( \frac{4\pi^2 c v^4}{3\kappa M} ut^4 \right) = \epsilon - \left( \frac{4\pi v^3}{3M} \right) \left( t^3 \frac{du}{dt} + 3ut^2 \right), \quad (13)$$

and finally we obtain eq. (9) in LP

$$t^3 \frac{du}{dt} + 3ut^2 = \left( \frac{3M}{4\pi v^3} \right) \epsilon - \left( \frac{\pi c v}{\kappa M} \right) ut^4. \quad (14)$$

## References

- [1] Li-Xin Li and Bohdan Paczyński. Transient events from neutron star mergers. *The Astrophysical Journal*, 507(1):L59, 1998.